

PC 3. Modeling gewydzid 40IN³/48, Co7H²/100.



EP
12768

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: G01N 33/16	A1	(11) International Publication Number: WO 79/00880 (43) International Publication Date: 1 November 1979 (01.11.79)
(21) International Application Number: PCT/US79/00210 (22) International Filing Date: 4 April 1979 (04.04.79) (31) Priority Application Number: 893,524 (32) Priority Date: 4 April 1978 (04.04.78) (33) Priority Country: US (71) Applicant: ALLEN, Robert, H.; 4001 S. Dexter, Englewood, CO 80110, United States of America. (72) Inventor: Applicant is also the inventor. (74) Agent: MARGOLIS, Donald, W.; 26 Garden Center, Suite 4, Broomfield, CO 80020, United States of America.		(81) Designated States: DE (European patent), FR (European patent), GB (European patent), JP. Published with: <i>International search report</i>
(54) Title: QUANTITATIVE TESTING FOR VITAMIN B12 (57) Abstract The vitamin B12 (cobalamin) level of human blood and mammalian tissue is routinely assayed utilizing radioisotope dilution assay (RIDA) techniques. The presence of vitamin "B12 analogues" in such samples, which analogues are erroneously assayed as true vitamin B12 utilizing prior art RIDA techniques, has been determined. Such errors are due to the fact that binding proteins commonly used in prior art RIDA techniques normally include proteins which bind both true vitamin B12 and previously unrecognized vitamin B12 analogues. The errors caused by the B12 analogues are avoided by using a composition for binding vitamin B12 which is substantially free of substances which bind vitamin B12 analogues. Moreover, by using both types of binders in separate assays, the amount of analogue present may be arrived at by taking the difference of the assays.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT

AT	Austria	LU	Luxembourg
BR	Brazil	MC	Monaco
CF	Central African Empire	MG	Madagascar
CG	Congo	MW	Malawi
CH	Switzerland	NL	Netherlands
CM	Cameroon	RO	Roumania
DE	Germany, Federal Republic of	SE	Sweden
DK	Denmark	SN	Senegal
FR	France	SU	Soviet Union
GA	Gabon	TD	Chad
GB	United Kingdom	TG	Togo
JP	Japan	US	United States of America

QUANTITATIVE TESTING FOR VITAMIN B₁₂BACKGROUND OF THE INVENTIONField of the Invention

The present invention relates to methods and materials for assaying mammalian blood and tissue. More specifically it relates to methods and materials for determining the amounts of vitamin B₁₂ and vitamin B₁₂ analogues in human plasma.

Prior Art

For many years it has been recognized that the assay of the vitamin B₁₂ level in humans is a valuable technique for diagnosing and subsequently treating certain diseases, such as for example, pernicious anaemia, post gastrectomy states, nutritional deficiencies, intestinal disorders, and others. Initially, vitamin B₁₂ was assayed microbiologically using either Euglena gracilis or Lactobacillus leichmannii. More recently, radioisotope dilution (RID) assays for B₁₂ have been utilized. Such radioisotope dilution assay techniques are well documented in the literature, see for example Lau, et.al. (1965) "Measurement of Serum B₁₂ Levels Using Radioisotope Dilution and Coated Charcoal," BLOOD, 26, 202, as modified by Raven et.al. (1968) "The Effect of Cyanide Serum and Other Factors on the Assay of Vitamin B₁₂ by Radio-Isotope Method Using ⁵⁷Co-B₁₂, Intrinsic Factor and Coated Charcoal," GUYS HOSPITAL REPORTS, 117, 89; and (1969) "Improved Method for Measuring Vitamin B₁₂ in Serum Using Intrinsic Factor, ⁵⁷Co-B₁₂ and Coated Charcoal," JOURNAL OF CLINICAL PATHOLOGY, 22, 205.

Such prior art radioisotope dilution assay of vitamin B₁₂ generally includes the steps of freeing the endogenous B₁₂ from its natural binding protein by boiling at a selected pH and then adding a measured amount of radioisotope ⁵⁷Co-B₁₂ and a limited amount of binding protein. All of the binding protein will be bound by some form of B₁₂ since the amount of radioisotope B₁₂ added is, by itself, sufficient to bind the small amount of protein. As both the natural B₁₂ and the radioactive B₁₂ compete to bind with the protein, the degree to which the radioactive count of the protein bound B₁₂ was



inhibited was thought to be indicative of the amount of natural B₁₂ present in the sample undergoing testing.

More specifically, in the technique of Lau et.al. as modified by Raven et.al., serum B₁₂ is separated from binding protein in the plasma sample by boiling with 0.25N HCl. Radioisotope B₁₂ is added to the reaction mixture and the B₁₂ mixture is reacted with protein, normally in the form of a commercially available binder. Then the free or unbound B₁₂ is separated from the protein bound B₁₂ by protein-coated charcoal and the radioactivity of the supernatant liquid containing the mixture of bound radioactive B₁₂ and bound non-radioactive B₁₂ counted for radioactivity. The serum B₁₂ concentration is then calculated from the count, often by comparison with a standard chart. Almost as soon as this technique began to be utilized it was recognized that the vitamin B₁₂ measurements it provided were usually inconsistent with the results obtained by other measuring techniques for B₁₂, such as the microbiological assays. Most often, the vitamin B₁₂ assay obtained by radioisotope dilution techniques have been found to be high. Many theories have been advanced to explain the cause of the high vitamin B₁₂ readings. However, it is believed that nowhere in the prior art is it recognized that there are substances in mammalian blood and tissue which react with certain non-specific protein binders in the radioisotope dilution assay technique to provide an analysis of vitamin B₁₂ which is apparently higher than the amount of B₁₂ actually in the sample. Additionally, it is believed that nowhere in the prior art is it recognized that most common and commercial RID assay protein binders are not specific to vitamin B₁₂, but that they are also capable of binding with the heretofore unknown B₁₂ analogues and thus provide erroneous B₁₂ assays.

BRIEF DESCRIPTION OF THE INVENTION

As has already been indicated, in the standard radioisotope binding assay for vitamin B₁₂, a known amount of radioactive vitamin B₁₂ is mixed with a prepared to-be-tested sample. Then, a known, but extremely limited, amount



of protein which is capable of binding with both the natural and radioactive vitamin B₁₂ is added to the mixture. Then, utilizing well known techniques, the radioactivity of the bound sample is compared, for example, with a standard curve to determine the amount of natural vitamin B₁₂ present in the tested sample. Such standard curves are initially established for use in RID assay, for example, by measuring the amount of bound radioactive B₁₂ in the presence of the same type and amount of protein binder, but with several different amounts of known non-radioactive B₁₂.

It has now been discovered, for what is believed to be the first time, that mammalian blood and tissue contain materials other than vitamin B₁₂ which couple with certain binding proteins which are commonly used in RID assays. For purposes of this specification and claims the non-vitamin B₁₂ materials which are capable of binding with such proteins will be herein referred to as "vitamin B₁₂ analogues," "B₁₂ analogues" or simply as "analogues." They are referred to as analogues, not due to their chemical structure, which is not known with certainty, nor in the commonly accepted chemical sense of the word "analogue." Rather they are referred to as analogues due to their reactivity with the binding proteins commonly used in RID assays. As will be shown in more detail, hereinafter, there are other similarities which have been discovered between vitamin B₁₂ and the newly discovered analogues which are present in mammalian blood and tissue.

After the presence of B₁₂ analogues was discovered it was then determined that protein binders commonly present in RID assays were: (1) Non-specific in binding to only vitamin B₁₂; and (2) reactive in binding with both vitamin B₁₂ and B₁₂ analogues; and (3) capable of reacting with both B₁₂ and B₁₂ analogues independent of pH. These are most commonly R proteins. Additionally, it has been determined that other protein binders, are: (1) Very specific in their reactivity substantially only with vitamin B₁₂; (2) substantially non-reactive with the B₁₂ analogues; and (3) non-reactive with either vitamin B₁₂ or B₁₂ analogues



in highly acid environments. These are most commonly proteins in the form of pure human intrinsic factor (IF), hog IF, rabbit IF, other IFs and vitamin B₁₂ specific binders.

5 In the past the problem has been that RID binders include substantial amounts of protein which is not specific to vitamin B₁₂. Therefore, the radioisotope dilution
10 assay utilizing that binder on samples which contain B₁₂ analogues will produce a measurement which indicates a greater amount of B₁₂ present in the plasma than exists in fact. As will be shown in more detail hereinafter, commercially available protein binders, which have heretofore been
15 labeled as containing intrinsic factor, in fact include only about 10% to about 30% intrinsic factor protein, while the balance of the protein in the binder is of a non-specific type, such as R protein. Thus, the protein materials in the commercial protein binders are capable of
20 indiscriminate reaction with the heretofore unrecognized vitamin B₁₂ analogue materials in mammalian blood and tissue. These extraneous reactions give RID analyses having the appearance of apparently higher vitamin B₁₂ content than the samples in fact contain. This is due to the fact that when
25 the binder includes protein which is non-specific to vitamin B₁₂ and which is capable of reacting with both vitamin B₁₂ and B₁₂ analogues, then the use of this protein in the radiobinding assay measures both the vitamin B₁₂ and the vitamin B₁₂ analogues which are present in the sample. However, in accordance with the present invention, when the proteins which are utilized are substantially specific to
30 vitamin B₁₂, such as substantially pure intrinsic factor, then in the RID assay one binds and measures substantially only the vitamin B₁₂ in the sample, without the measurement of extraneous B₁₂ analogues. This provides a more accurate vitamin B₁₂ RID assay.

35 Based on these discoveries it is proposed that in the practice of RID assay only protein which is specific in its reaction to vitamin B₁₂ be utilized. Alternatively, it is proposed that mixtures of vitamin B₁₂ specific and



non-specific binding proteins be treated, for example, with an excess of material which will bind or inactivate only the non-specific binding proteins, such as vitamin B₁₂ analogues, prior to its use in RID assays, so that the non-specific protein will be substantially unavailable for reaction with any vitamin B₁₂ or analogues in a sample when the RID assay is conducted. In yet another modification of the present invention, crude binder, including non-specific binding proteins, is subjected to proteolytic enzyme treatment prior to utilization as a vitamin B₁₂ binder in RID assays. Such proteolytic enzyme treatment destroys the binding ability of the non-specific proteins without destroying the binding ability of the proteins which are specific to vitamin B₁₂.

Utilizing the techniques of the present invention, the B₁₂ analogues can be assayed by analyzing the amount of vitamin B₁₂ present utilizing, for example, a vitamin B₁₂ specific binder, then assaying the sample utilizing a non-specific binder and determining the difference between the two assays as a measure of the amount of vitamin B₁₂ analogues present.

These and other techniques are readily determined, once, as taught for the first time by the present invention, the presence of B₁₂ analogues in mammalian blood and tissue is recognized.

The foregoing and other objects, features and advantages of the invention will be apparent from the following more particular description of preferred embodiments of the invention.

DETAILED DESCRIPTION OF THE INVENTION

In the following examples and tables certain chemical components were utilized. For ease of communication they have been given shortened names in the text. The concordance between the "component" names and their actual compositions is as follows:

<u>Components</u>	<u>Actual Composition</u>
A. Buffer	1.0M Tris (hydroxymethyl) aminomethane-HCl pH 10.0



6

- B. Albumin
C. Salt
D. Boiled buffer
- 5
- 10
- E. Standard
(100 pg/ml B₁₂)
- 15
- 20
- F. Standard
(1000 pg/ml B₁₂)
- 25 G. (⁵⁷Co) B₁₂
- H. Binder
- 30
- 35
- Bovine serum albumin, 2 mg
per ml in H₂O
0.15M NaCl
(1 part) 0.5M sodium acetate-
HCl pH 4.5
(1 part) 0.01M KPO₄ pH 7.5,
0.15M NaCl
(2 parts) 50 µg per ml KCN
in 0.15M NaCl
The complete solution is heated
of 45 min. at 100°C.
Solution D containing 100 pg
per ml vitamin B₁₂. The so-
lution is heated for 45 min.
at 100°C after the vitamin B₁₂
is added. The concentration
of vitamin B₁₂ in the stock
solution used to make component
E is determined by its light
absorbance at 278, 361 and 550
nm.
Same as component E except that
the vitamin B₁₂ concentration
is pg/ml.
1000 pg per ml of (⁵⁷Co) B₁₂,
(150-300 µCi/µg), in H₂O.
Present in 0.01M Tris-HCl pH
8.2, containing 0.15M NaCl and
50 µg per ml bovine serum albumin.
Binders are diluted in this
solution to reach a concentra-
tion of 700 pg per ml of vitamin
B₁₂ binding ability.
Individual binders are as follows:
1) Human intrinsic factor
(Human IF) -
Human gastric juice contain-
ing more than 95% intrinsic



7

factor based on assays employing inhibition of vitamin B₁₂ binding with anti-intrinsic factor antibodies (95% inhibition) and cobinamide (5% inhibition).

5

- 2) Human R protein (Human R)- Human saliva containing more than 95% R protein based on assays employing inhibition of vitamin B₁₂ binding with cobinamide (95% inhibition) and anti-intrinsic factor antibodies (5% inhibition)

10

15

- 3) Hog intrinsic factor (Hog IF) - This protein was purified from "Hog intrinsic factor concentrate" by affinity chromatography on vitamin B₁₂-Sephadex employing gradient elution with guanidine-HCl followed by gel filtration. The final preparation contained more than 95% intrinsic factor based on assays employing inhibition of vitamin B₁₂ binding with anti-intrinsic factor antibodies (95% inhibition) and cobinamide (5% inhibition).

20

25

30

- 4) Hog R protein (Hog R) (Also designated in the scientific literature as Hog non-intrinsic factor concentrate)

35



8

This protein was purified from "Hog intrinsic factor concentrate" as described above in 3). The final preparation contained more than 95% R protein based on assays employing inhibition of vitamin B₁₂ binding with anti-intrinsic factor antibodies (5% inhibition) and cobinamide (95% inhibition).

- 5) Rabbit intrinsic factor (Rabbit IF) - An extract of rabbit gastric mucosa containing more than 95% intrinsic factor based on assays employing inhibition of vitamin B₁₂ binding with anti-intrinsic factor antibodies (95% inhibition) and cobinamide (5% inhibition).
- 6) Hog intrinsic factor concentrate (Hog IFC) - A crude extract of hog pyloric mucosa. It contained 25% Hog IF and 75% Hog R based on assays employing inhibition of vitamin B₁₂ binding with anti-intrinsic factor antibodies (25% inhibition) and cobinamide (75% inhibition).
- 7) Hog IFC + Cobinamide - Hog IFC containing the vitamin B₁₂ analogue cobinamide



9

([CN, OH] Cbl) in a molar amount equal to 100 times the total vitamin B₁₂ binding ability, i.e. a 100 fold excess of cobinamide.

5

8) Hog IFC + CN-Cbl [bde-OH] - The same as item 7) above except that the analogue added is CN-Cbl [bde-OH] and is present in a 1000 fold molar excess.

10

9) Hog IFC + [3,5,6-Me₃BZA] (CN,OH)Cba - The same as item 7) above except that the analogue added is [3,5,6-Me₃BZA] (CN, OH)Cba.

15

10) Digested Hog IFC - Hog IFC incubated with bovine pancreatic trypsin (2 mg per ml) and bovine pancreatic chymotrypsin (2 mg per ml) for 60 min. at 37°C.

20

I. Charcoal

A solution containing 25 mg per ml neutral charcoal (Norit) and 5 mg per ml bovine serum albumin in H₂O.

25

J. unknown sample

Samples containing unbound vitamin B₁₂ are diluted in solution D (see above). Samples containing bound vitamin B₁₂, such as serum, are prepared as follows:

30

(1 part) sample

(1 part) 0.5M sodium acetate-HCl pH 4.5

35

(2 parts) 50 µg per ml KCN in 0.15M NaCl



10

The complete mixture is heated at 100°C for 45 min. followed by centrifugation at 5000x g at 4°C for 20 min. The supernatant is removed and used for assay.

5

Each of the RID assays referred to herein utilized the components referred to above. The method and order of utilizing the components is that set forth in Table 1. That is, components A, B, C, etc. or the buffer, albumin, and salt, respectively, etc. were added in the order, from left to right, shown in Table 1. After the addition of $^{57}\text{Co-B}_{12}$ the components are mixed thoroughly to mix both the naturally occurring B_{12} and the radioisotope B_{12} to make them compete and equally available to react with the binder. After the addition of the binder, H, the components were again mixed thoroughly, and then incubated for 45 minutes at about 37°C. Charcoal was then added to the incubated mixture and the components again mixed thoroughly and incubated for another 5 minutes at room temperature. This was followed by centrifuging at 2000 x g at 4°C for 30 minutes. Then 1000 μl of the resulting supernatant liquid is pipetted from the sample and a determination of the amount of $^{57}\text{Co-B}_{12}$ present is made. The amount of $^{57}\text{Co-B}_{12}$ is indicative of the amount of natural B_{12} in the tested sample, with lesser amounts of $^{57}\text{Co-B}_{12}$ being indicative of greater amounts of natural vitamin B_{12} in the sample.

15

20

25

Calculations of vitamin B_{12} , utilizing the data obtained in the foregoing manner, is made as follows:

30

Calculation of data from radiobinding assay for plasma vitamin B_{12} assay as outlined in Table I

35

- 1) The values in tubes 3 and 4, the "blank" tubes without binder are averaged and subtracted from all other tubes starting with tube 5.
- 2) The background radiation is subtracted from tubes 1 and 2 and these values are averaged.
- 3) Tubes 5 and 6 are averaged. This value should be at



Table I

Flow Sheet for Radiobinding Assay for Vitamin B12

Tube #	A Buffer (ul)	B Albumin (ul)	C Salt (ul)	D Boiled Buffer (ul)	E Standard 100pg/ml B12 (ul)	F Standard 1000pg/ml B12 (ul)	J Unknown Sample (ul)	G [57Co] B12 (ul)	H Binder (ul)	I Charcoal (ul)
1,2	225	50	875	800	-	-	-	50	-	-
3,4	225	50	375	800	-	-	-	50	-	500
5,6	225	50	325	800	-	-	-	50	50	500
7	225	50	325	720	80	-	-	50	50	500
8	225	50	325	640	160	-	-	50	50	500
9	225	50	325	560	240	-	-	50	50	500
10	225	50	325	300	500	-	-	50	50	500
11	225	50	325	720	-	80	-	50	50	500
12	225	50	325	640	-	160	-	50	50	500
13	225	50	325	450	-	350	-	50	50	500
14	225	50	325	-	-	800	-	50	50	500
Unknown Sample	225	50	325	-	-	-	800	50	50	500



12

least 15% below the average value for tubes 1 and 2 to insure that all of the binder is saturated in the presence of (^{57}Co) B_{12} alone.

- 4) Values for each tube beginning with tube 7 are divided by the average of tubes 5 and 6 to give values for "% trace binding."
- 5) Percent trace binding for tubes 7-14 are used to obtain a standard curve. We plot % trace binding on the ordinate of logit-log paper versus pg vitamin B_{12} on the log scale.
- 6) The amount of vitamin B_{12} in unknown samples is determined by interpolation from the standard curve or data of % trace binding versus pg vitamin B_{12} .
- 7) The standard curves for all of the various binders used are virtually indistinguishable and vary little from day to day. Nevertheless, a complete standard curve is always obtained for every binder with each set of assays. Representative data obtained with the assays are present in Table II.

20 Evidence as to the Origin and Existence of Vitamin B_{12}
Analogues in Mammalian Blood and Tissue

Once the problem of the prior art is recognized, that is; that there are vitamin B_{12} analogues present in mammalian blood and tissue, it becomes a relatively simple matter to prove the existence and chemistry of such analogues. It is also appropriate to prove that the various steps of the RID assay do not cause the B_{12} analogues to be formed, for example, from vitamin B_{12} .

In one instance this has been most convincingly shown by obtaining pure crystalline vitamin B_{12} , subjecting various known concentrations of it to the same conditions used to extract endogenous vitamin B_{12} from blood and tissue samples (boiling for 45 minutes in the same extraction solution) and then analyzing them by RID assay using several binding proteins, for example, in the form of human IF, hog IF, human R, hog R and hog IFC on different portions of the same extracted vitamin B_{12} samples..



Table II

Standard Curves obtained with the radiobinding assay for vitamin B₁₂ using various binders

Vitamin B ₁₂ added (a) (pg)	Z trace binding observed with various binders (b)			
	Human IF	Human R	Hog IF	Hog R
0	(100)	(100)	(100)	(100)
8	90	93	94	89
16	85	84	83	80
24	79	78	74	76
50	64	60	58	59
80	45	44	44	43
160	30	28	28	28
350	15	14	15	14
800	6	6	7	6

(a) The vitamin B₁₂ was boiled for 45 min. at 100°C in the same solution used to extract endogenous vitamin B₁₂ from human plasma.

(b) Assays were performed on different days.



Referring to Table II, it will be seen that when various known amounts of pure vitamin B₁₂, ranging from about 8 pg to about 800 pg were tested with various protein binders, that in each instance, the percent of radioactive trace binding, or more accurately, the inhibition of ⁵⁷Co-B₁₂ binding, observed was substantially the same for each binder. It is thus seen, that regardless of which protein binder is utilized, the percent binding, i.e. inhibition of the (⁵⁷Co)-B₁₂ is substantially the same. This is indicative of the fact that during preparation for RID assay the pure vitamin B₁₂ was not converted to analogues of the type which have now been observed in mammalian blood and tissue. It is also indicative of the fact that in the absence of interfering masking components in the samples, such as B₁₂ analogues, any of the binding proteins can be utilized to provide substantially equally accurate RID assays of vitamin B₁₂. Furthermore, the data in Table II should be suitable as a standard in the determination of vitamin B₁₂ by the same RID assay.

By comparison, when endogenous vitamin B₁₂ was extracted from serum from 74 normal blood donors (37 women, 37 men, ages 17-61) and tested utilizing the same binding proteins with the exception of hog IFC which was not used, the results were quite different. In every case in which serum from normal donors was tested greater inhibition of ⁵⁷Co-B₁₂, and therefore greater apparent vitamin B₁₂, was observed with assays employing, as the binder, human R or hog R than was observed with assays employing human IF or hog IF. The data on the 74 normal donors is included in Table III. Other data concerning patients with diagnosed vitamin B₁₂ deficiencies are present, and comparisons between the normal donors and patients have also been made on Table III, and will be discussed in more detail hereinafter. Referring to Table III it is seen that the mean endogenous vitamin B₁₂ RID assay levels, in terms of pg of vitamin B₁₂ per ml of serum, are 548 and 542 for human R and hog R,



Table III

Distribution of serum vitamin B₁₂ values as measured with various binders for 74 normal subjects and 71 patients with clinical evidence of vitamin B₁₂ deficiency
All assays were performed at pH 9.0

Serum Vitamin B ₁₂ (pg/ml)	Human R		Hog R		Human IF		Hog IF		Human R minus Human IF		Hog R minus Hog IF	
	Normals	Patients	Normals	Patients	Normals	Patients	Normals	Patients	Normals	Patients	Normals	Patients
0-24						8		11				
25-49						6		5				
50-74						5		3		1		1
75-99		1		1		2		2		2		3
100-124		2		1					2	3		3
125-149		2		5	4		1		7	3		3
150-174		3		1	5		3		6	5		5
175-199		4		3	2		3		6	3		3
200-224	1	1		2	5		4		8	2	11	2
225-249		3	1	1	6		4		2	3	6	3
250-274				3	10		9		10		4	
275-299		1	2	1	4		3		5		1	
300-324	1	2	4	2	12		4		6		3	
325-349	3	1	2	1	5		6		4	1	5	1
350-374	5	1	1				10		4		1	
375-399	3		2		3		4		4		2	
400-424	5		3		4		5		5		1	
425-449	3		4		3		2		2			
450-474	4		6		2		3				1	
475-499	5		5				3				1	
500-524	7		4		1		3					
525-549	3		4		1							
550-574	5		3		2		1				1	
575-599	1		2		1				1			
600-624	2		4		1		3				1	
625-649	5		4									
675-699	2		3									
700-724	3		6				1					
725-749	2											
750-774	3		1				1					
775-799			1		1							
800-824			2									
825-849												
850-874	1		1									
875-899	1		2				1					
900-924	1		1									
925-949	2		1									
950-974	1											
975-999	1											
1000-1024												
1025-1049												
1050-1074	1											
1075-1099												
1100-1124												
1125-1149			1									
1150-1174												
1175-1199												
1200-1224												
1225-1249	1											

range of serum B ₁₂	220-1230	85-355	243-1135	84-342	130-785	0-78	135-880	0-86	70-375	56-337	50-605	72-332
mean of serum B ₁₂ (a)	348		542		298	36	361	33	254	169	212	171
mean ± 2 Std. Dev.	282- 1040		276- 1065		136- 656		157- 717		70- 661		58- 531	

(a) Based on values obtained with log vitamin B₁₂. Log values were used since these gave a normal curve while the curve using untransformed values were skewed to the right.



respectively, but only 298 and 361 for human IF and hog IF, respectively. This demonstrates that something is present in extracts of normal human serum which inhibits the vitamin $^{57}\text{Co-B}_{12}$ binding ability of human R and hog R to a greater extent than it inhibits the binding ability of human IF and hog IF. Under current RID assay techniques the greater inhibition which is found using human R and hog R is analyzed to indicate a higher vitamin B_{12} content. It is those substances, which have now been found to be present in human blood serum and which preferentially inhibit $^{57}\text{Co-B}_{12}$ binding of human R and hog R, which have been herein denominated as "vitamin B_{12} analogues."

Chemical Nature and Properties of Vitamin B_{12} Analogues

The vitamin B_{12} analogues, which are herein for the first time identified as being present in mammalian blood and tissue, have been isolated by paper chromatography and compared with pure vitamin B_{12} . Vitamin B_{12} and the so-called "vitamin B_{12} analogues" were found to have the following properties in common: (1) Both were adsorbed to charcoal and remained adsorbed when the charcoal was washed with 5% phenol; (2) Both were eluted from charcoal when the charcoal was washed with 67% acetone; (3) Both were extracted from aqueous solution into phenol and remained in the phenol phase even when the phenol was washed repeatedly with water; (4) Both passed into the aqueous phase when the phenol layer was dissolved in an excess of diethyl ether; (5) Both eluted with similar apparent molecular weights (approximately 1356) during gel filtration on columns of Bio-Rad P-4 polyacrylamide; (6) Both were adsorbed to a column of Sepharose-2B agarose that contained covalently bound hog R protein and both remained bound when the column was washed with 0.1M glycine-NaOH pH 10.0, 1.0M NaCl, and both were eluted from the Sepharose with either 85% phenol or 60% pyridine. Because of these similarities the newly discovered material is seen to be similar to vitamin B_{12} and is thus referred to as vitamin B_{12} analogue.



The chemical nature and structure of the newly discovered vitamin B₁₂ analogues which are now found to be present in mammalian blood and tissue is not known. An effort was made to compare them with chemically true forms of vitamin B₁₂, sometimes referred to in the literature as analogues of vitamin B₁₂, namely CN-B₁₂, OH-B₁₂, adenosyl-B₁₂ and CH₃-B₁₂, already known to be present in serum and tissues. This was done by adding 500 pg of each of these four known forms of vitamin B₁₂ to four different portions of the same human serum, in the dark. Prior to the additions the serum contained 250 pg and 450 pg of vitamin B₁₂ as assayed by RID using human IF and human R, respectively, thus exhibiting a difference of 200 pg. After addition of the materials to the serum, each was allowed to incubate in the dark for 15 minutes to allow binding of the added known forms of vitamin B₁₂ to the binding proteins normally present in the serum. Then the serum, with the added forms of vitamin B₁₂ was extracted utilizing standard conditions and the apparent amount of vitamin B₁₂ assayed by RID utilizing both human R protein and human IF protein. Both the human R and human IF assays showed an increase in the apparent amount of vitamin B₁₂ of about 500 pg. However, the original difference observed between the values obtained with the human R protein and the human IF protein, i.e. 200 pg, did not change. If any of the added known forms of vitamin B₁₂ in the human serum had been converted to the newly discovered analogues, then the assays would have shown an increase in the difference. This provides evidence that the newly discovered vitamin B₁₂ analogues were not formed from any of the known endogenous forms of native vitamin B₁₂ during the extraction procedure.

Isolation of Vitamin B₁₂ Analogue

The materials which are herein designated as "vitamin B₁₂ analogues" and which have been found to preferentially inhibit R proteins in the vitamin B₁₂ assays were substantially separated from endogenous vitamin B₁₂ by the following purification scheme. A trace amount of 150 pg ⁵⁷Co-B₁₂, was added to 1800 ml of freshly collected normal human



plasma. The added $^{57}\text{Co-B}_{12}$ was sufficiently small that it did not interfere with subsequent RID assays. After incubating at room temperature for 30 minutes the vitamin B_{12} was extracted and assayed under standard conditions.

5 When human IF binder was utilized in the RID assay the extract was found to contain 1050 ng of vitamin B_{12} , but when human R protein was utilized as the binder it appeared to contain 2030 ng of vitamin B_{12} , almost twice as much vitamin B_{12} . The extract was then passed through a column of

10 Sepharose containing covalently bound hog R protein. The column retained greater than 99% of the $^{57}\text{Co-B}_{12}$ as well as the endogenous vitamin B_{12} as assayed by RID with human IF or human R protein. After the column was washed with a variety of buffers and water the material was eluted with

15 60% pyridine, taken to dryness under vacuum, dissolved in water, and adsorbed onto charcoal. The charcoal was washed with 5% phenol followed by water and the mixture of vitamin B_{12} , $^{57}\text{Co-B}_{12}$ and analogue B_{12} was eluted from the charcoal with 67% acetone. The material was again taken

20 to dryness under vacuum, dissolved in water, and then separated utilizing 19 inch long Whatman 3MM paper for paper chromatography and a solvent system consisting of 800 ml sec-butanol, 8 ml glacial acetic acid, 6 mmol HCN and 400 ml water. The chromatography was performed in the descend-

25 ing manner for 30 hours at room temperature in an environment that inhibited evaporation of the solvent. The paper chromatogram was allowed to dry in a fume oven and divided into 38 one-half inch fractions and numbered, with fraction 1 starting at the point of application and number 38 being

30 at the lowest point on the chromatogram. Each one-half inch fraction was then incubated with 5 ml of water at 4°C for twelve hours to elute the vitamin B_{12} , $^{57}\text{Co-B}_{12}$ and B_{12} analogues. The water was then removed and taken to dryness under vacuum. Each dried fraction was then dissolved

35 in 2.5ml of water and assayed for $^{57}\text{Co-B}_{12}$ and for vitamin B_{12} using a variety of binding proteins. The final recovery of $^{57}\text{Co-B}_{12}$ was 64%. The apparent recoveries of vitamin B_{12} were 75% when using human IF in the assay

and 66% when using human R in the assay. The results of the assays employing the 38 fractions obtained by paper chromatography are presented in Table IVA, IVB, and IVC. Similar data concerning paper chromatography of $^{57}\text{Co-B}_{12}$ and pure vitamin B_{12} , for reference as a control, are presented in Table IVD. The data in these several chromatography tables summarized for convenience in Table IVE reveals that the behavior of $^{57}\text{Co-B}_{12}$ that was extracted from human plasma did not change its chromatographic behavior, and thus was not altered during the standard extraction procedure or any of the purification steps. In a similar manner it is postulated that true vitamin B_{12} is not altered in any of the purification or process steps of the assay.

Referring to the control chromatogram of Table IVD, it is seen that the several RID assays of pure vitamin B_{12} performed variously with human R protein, human IF protein and hog IFC gave substantially a single symmetrical peak of activity. In each instance greater than 95% of the vitamin B_{12} was found to be present in fraction 14 through 16. Similar results, as shown in Table IVB, were obtained from the paper chromatogram of the plasma extract when the binding protein was human IF, hog IF and rabbit IF. These data are an indication that these three IF binding proteins are substantially specific in their binding ability to vitamin B_{12} , and substantially non-reactive with vitamin B_{12} analogues present in plasma.

Efforts were made to modify hog IFC, which is a commonly used binder in RID assays and which has been found to contain as much as 90% hog R protein and as little as 10% hog IF, by removing or inactivating the hog R. In several instances the hog IFC was incubated with an excess amount of three chemically synthesized vitamin B_{12} analogues before it was utilized in the RID assay. Referring to Table IVC, it is seen that after this modification the chromatogram results obtained utilizing the modified hog IFC closely resemble the results obtained with substantially pure hog IF. It is therefore seen, that in the practice of the

Table IVA
[57Co] B12 and plasma B12 assayed after elution from paper chromatography.
All assays were performed at pH 9.0 except where indicated.

B12 assayed with various binders

Fraction #	[57Co] B12 (cpm)	Human R		pH 1.8 Human R		Hog R		pH 1.8 Hog R		Hog IFC	
		(ng)	(%)	(ng)	(%)	(ng)	(%)	(ng)	(%)	(ng)	(%)
1		2		1		1		1		1	
2											
3											
4		1		1		1		2		1	
5		2				2		3		2	
6		3		3		2		4		2	
7		4		4		4		5		3	
8		4		3		4		5		3	
9		7		6		6		8		5	
10	300	24		29		26		29		16	
11	500	42		54		36		49		26	
12	300	25		34		26		38		19	
13	200	31		44		30		55		22	
14	4600	209		235		176		240		180	
15	14000	562		615		461		576		474	
16	2100	103		122		84		122		91	
17	300	85		118		85		128		53	
18	300	68		68		68		100		45	
19	200	23		17		17		27		15	
20		45		39		39		60		24	
21		42		25		25		52		21	
22		8		5		7		8		5	
23		8		4		5		8		4	
24		7		4		5		7		3	
25		9		1		6		4		4	
26		16		1		10		2		5	
27		5		1		3		1		2	
28		2		1		1		1		1	
29		4				2		1		2	
30		1				1		1		1	
31		1									
32-38											

Table IVB

Assay of [⁵⁷Co] B₁₂ and plasma B₁₂ after elution from paper chromatography.
All assays were performed at pH 9.0.

Fraction #	[⁵⁷ Co] B ₁₂		B ₁₂ assayed with various binders		Rabbit IF	
	(cpm)	(%)	Human IF (ng)	Hog IF (ng)	(ng)	(%)
1			1		1	
2						
3						
4					1	
5				1		
6				1		
7				1		
8				1		
9			1			
10	300	1	4	5	8	
11	500	2	7	10	16	
12	300	1	5	7	10	
13	200	1	14	10	13	
14	4600	20	169	169	170	
15	1400	61	488	450	480	
16	2100	9	73	72	87	
17	300	1	17	25	42	
18	300	1	11	15	25	
19	200	1	3	5	5	
20			5	7	8	
21			2	4	4	
22-38						



Table

Assay of [⁵⁷Co] B12 and plasma B12 after elution from paper chromatography.
All assays were performed at pH 9.0.

Fraction #	[⁵⁷ Co] B12		Hog IFC		Hog IFC + Cobinamide		Hog IFC + CN-Chl[bde-OH]		Hog IFC + [3,5,6-Me ₃ BZA] (CN, OH)Cba		Hog IFC Digested	
	(cpm)	(%)	(ng)	(%)	(ng)	(%)	(ng)	(%)	(ng)	(%)	(ng)	(%)
1			1								1	
2												
3			1				1				1	
4			2				1				1	
5			2				1				1	
6			3				1				2	
7			3				1				2	
8			5				1				3	
9			16		1		4				9	
10	300	1	26	2	10	1	10	1			15	1
11	500	2	19	3	8	1	8	1			11	2
12	300	1	22	2	9	1	10	1			14	2
13	200	1	180	17	174	21	170	22			153	19
14	4600	20	474	46	474	58	435	55			402	50
15	14000	61	91	9	83	10	81	10			84	10
16	2100	9	53	5	25	3	27	3			36	4
17	300	1	45	4	16	2	19	2			27	3
18	300	1	15	1	4		4	1			9	1
19	200	1	24	2	7	1	7	1			13	2
20			21	2	3		4	1			9	1
21			5	1				1			2	
22			4								1	
23			3								1	
24			4								1	
25			5								2	
26			2								1	
27			1								1	
28			2								1	
29			1								1	
30			2								2	
31-38			1								1	

BUREAU

COMP

Table IVE

Summary of the data in Tables IVA & IVD involving assays of [⁵⁷Co] B₁₂ and B₁₂ after elution from paper chromatography. Assays for B₁₂ were performed at pH 9.0 except where indicated.

Sample Chromatographed	Item Assayed	Assay Binder	Chromatogram Fractions							
			1-13		14-16		17-38		1-38	
			(ng)	(%)	(ng)	(%)	(ng)	(%)	(ng)	(%)
[⁵⁷ Co] B ₁₂ + B ₁₂	[⁵⁷ Co] B ₁₂	-	-	5	-	92	-	2	-	100
"	B ₁₂	Human R	7	2	336	97	5	1	348	100
"	B ₁₂	Human IF	5	1	322	98	2	1	329	100
"	B ₁₂	Hog IFC	11	3	319	95	7	2	337	100
[⁵⁷ Co] B ₁₂ + plasma B ₁₂	[⁵⁷ Co] B ₁₂	-	-	6	-	91	-	3	-	100
"	B ₁₂	Human R	145	11	874	65	324	24	1343	100
"	B ₁₂	Human R (pH 1.8)	179	12	972	68	283	20	1434	100
"	B ₁₂	Hog R	138	12	721	64	274	24	1133	100
"	B ₁₂	Hog R (pH 1.8)	199	13	938	61	402	26	1539	100
"	B ₁₂	Hog IFC	100	10	745	72	185	18	1030	100
"	B ₁₂	Human IF	31	4	730	91	38	5	799	100
"	B ₁₂	Hog IF	37	5	691	88	56	7	784	100
"	B ₁₂	Rabbit IF	53	6	737	84	84	10	874	100
"	B ₁₂	Hog IFC + Cobinamide	32	4	731	89	54	7	818	100
"	B ₁₂	Hog IFC + CN-Cbl [bde-OH]	47	6	624	83	81	11	752	100
"	B ₁₂	Hog IFC + [3,5,6-Me ₃ BZA] (CN, OH) Cba	38	5	686	87	61	8	785	100
"	B ₁₂	Hog IFC - Digested	60	7	639	80	102	13	801	100



present invention, mixtures of protein including both vitamin B₁₂ specific binding protein and binding protein which is not specific to vitamin B₁₂ can be modified by the addition of an excess amount of vitamin B₁₂ analogue, 5 by which process the analogue binds with the non-specific protein to render it substantially bound or inactive so that it is not available to react with vitamin B₁₂ or (⁵⁷Co) B₁₂ present in samples undergoing RID tests. The amount of vitamin B₁₂ analogue to be added to a mixture of 10 specific and non-specific proteins in order to bind or inactivate the non-specific proteins may vary over a wide range, depending on both the proteins which are present and the vitamin B₁₂ analogues which are utilized as the binding or inactivating material. Generally speaking, for the exam- 15 ples shown in Table IVC, cobinamide may be added in an amount equal to that required for complete binding of the non-specific protein, up to an amount as much as ten million times greater than the amount needed to bind the protein, with the preferred range being about ten to about ten 20 thousand times in excess of that required for complete binding. CN-Cb (bde-OH), known as CoB-cyano-cobamic a,c,g-triamide may be utilized in an amount at least about ten times to about ten million times in excess of the amount required to bind with the non-specific protein, with an 25 amount of about one hundred to about one hundred thousand times excess being preferred. The (3,5,6-Me₃BZA) (CN,OH)-cba, known as Co (3,5,6-trimethylbenzimidazole) cobamide should also be utilized in amounts from about one to ten million times in excess of the amount of non- 30 specific protein, with an amount in the range of about ten to about ten thousand times excess being preferred. Suitable amounts of other vitamin B₁₂ analogues may be utilized in a similar manner to bind or inactivate non-specific proteins present in mixtures with specific proteins in order 35 to obtain a preparation of binding protein which will substantially bind only ⁵⁷Co-B₁₂, or the vitamin B₁₂ naturally present in the samples being tested, and thus



give a more accurate quantitative RID assay of vitamin B₁₂ in samples undergoing tests.

Again, referring to Table IVC, data on samples of hog IFC digested with trypsin and chymotrypsin are shown. These 5 and other proteolytic enzymes are specific in their ability to substantially digest R proteins while leaving intrinsic factor proteins unaffected and available as substantially the only protein for binding ⁵⁷Co-B₁₂ and vitamin B₁₂ in RID assays. Other enzymes, including, for example, elastase 10 may be utilized for the same purpose. The amount of the enzymes utilized is in the range of about 0.01 to about 100 milligrams per milliliter of protein treated, with a preferred amount being about 0.05 to about 40 milligrams per milliliter of protein. Utilizing this proteolytic enzyme 15 digestion process a protein binder is provided which substantially binds only ⁵⁷Co-B₁₂ and vitamin B₁₂ and is not affected by vitamin B₁₂ analogues in the samples being tested and which therefore gives a more accurate RID assay than is obtained when utilizing the original mixture of hog 20 IFC proteins which included non-specific proteins which would have been capable of reacting with the newly discovered vitamin B₁₂ analogues in samples to give inaccurate assays as to the amount of vitamin B₁₂ in test samples.

Now, referring to Table IVA, when samples from the 25 plasma extract chromatogram were assayed for vitamin B₁₂ with human R, hog R and hog IFC, different results were obtained than when those samples were assayed with human IF, hog IF, rabbit IF or hog IFC treated with vitamin B₁₂ analogues or hog IFC digested with proteolytic enzymes. 30 In each case where human R, hog R or untreated hog IFC were utilized as the binding protein the tests gave the appearance that more vitamin B₁₂ was present in the chromatogram samples, especially in fraction 1 through 13 and 17 through 38. This observation, when taken with the above data, provides strong evidence that normal human plasma contains a 35 number of vitamin B₁₂ analogues that compete with ⁵⁷Co-B₁₂, in significant amounts, for binding to R protein. It also indicates that such activity on the part of the B₁₂



analogues is substantially absent when the binding protein utilized in the RID assay is substantially specific to vitamin B₁₂.

It should also be noted, see Table IVA, that the chromatogram data suggests that the lack of specificity of human R and hog R is unchanged when RID assays are performed at acid pH. This indicates that erroneous results will be obtained for the true vitamin B₁₂ content of samples which contain vitamin B₁₂ analogues when RID assays are performed at acid pH.

Using the same techniques and criteria described above it has been discovered that vitamin B₁₂ analogues are not only present in serum obtained from human blood, but are also present in mammalian tissues in even higher concentrations than they are in blood. Vitamin B₁₂ analogues extracted from mammalian tissues have been purified using the same schemes as described above. When analyzed utilizing paper chromatography, they exhibited similar mobilities to those of the vitamin B₁₂ analogues observed in the samples extracted from blood serum. Since larger amounts of the vitamin B₁₂ analogues are present in tissue, they can be observed visually as red or orange spots during paper chromatography. The absorption spectra of vitamin B₁₂ analogues purified from tissue extracts have been determined and demonstrate that they are similar to, but distinct from, the absorption spectrum of true vitamin B₁₂. These observations provide additional evidence that the materials in blood serum which preferentially react with R proteins and not intrinsic factor proteins are in fact varieties of vitamin B₁₂ analogues.

The newly discovered vitamin B₁₂ analogues also differ from vitamin B₁₂ in terms of their biological activity. Thus, as shown in Table V, the serum vitamin B₁₂ values obtained with Euglena gracilis for eleven patients diagnosed to have vitamin B₁₂ deficiency were substantially similar to the results obtained by RID assay using human IF or hog IF as the binding protein. It is to be further noted, that all of the values obtained by either microbiologic assay or



Table V

Vitamin B₁₂ levels in 21 patients with clinical evidence of vitamin B₁₂ deficiency

Patient	Vitamin B ₁₂ assayed with various binders				Vitamin B ₁₂ assayed with <i>Euglena</i> <i>Gracilis</i>
	Human R (pg/ml)	Hog R (pg/ml)	Human IF (pg/ml)	Hog IF (pg/ml)	
1	155	138	16	0	43
2	310	295	51	86	52
3	310	255	24	35	46
4	135	132	0	22	40
5	215	250	52	60	25
6	347	342	0	0	56
7	102	120	0	0	35
8	240	242	58	65	79
9	160	125	38	23	0
10	85	84	0	12	23
11	235	255	5	11	0
12	188	190	41	38	
13	178	192	48	38	
14	298	305	53	57	
15	355	310	78	82	
16	178	210	48	22	
17	128	140	31	8	
18	106	190	50	40	
19	163	155	0	0	
20	178	132	25	0	
21	230	215	75	41	
Mean (1-11)	209	203	22	29	36
Mean (1-21)	205	204	33	30	
Normal Range (mean \pm 2 std. dev.)	220-1230	245-1135	136-656	157-717	(>130)
Number within normal range	9 (43%)	10 (48%)	0	0	0



by assay using human IF or hog IF were substantially lower than the values obtained when the RID assay was carried out utilizing human R or hog R as the binding protein.

This indicates that the vitamin B₁₂ analogues which have now
5 been identified in mammalian blood and tissue do not possess vitamin B₁₂ activity of the type which is required to promote the growth of Euglena gracilis.

Data was obtained on ten additional patients diagnosed to be vitamin B₁₂ deficient and the total of 21 patients
10 with vitamin B₁₂ deficiency are shown in Table V. In each of the 21 patients the vitamin B₁₂ values found when the RID assay was carried out utilizing human IF or hog IF were below the range of vitamin B₁₂ values found in a control group of 74 normal subjects. However, when the RID assay
15 was carried out utilizing human R or hog R only about half of the 21 vitamin B₁₂ deficient patients were found to assay below the range of normal subjects for vitamin B₁₂. This indicates that where the newly found vitamin B₁₂ analogues are present in the samples being tested, and the
20 binding protein is not specific to vitamin B₁₂, the resulting assays may suggest that a truly vitamin B₁₂ deficient patient is not within the deficient range. This may lead to delay of treatment of that patient for vitamin B₁₂ deficiency. It also indicates that the vitamin B₁₂ ana-
25 logues that have now been discovered lack the therapeutic or beneficial activity of vitamin B₁₂ in the sense of being unable to prevent the hematologic and/or neurologic abnormalities associated with vitamin B₁₂ deficiency.

30 There are many commercial RID assay type kits available for the assay of vitamin B₁₂ in clinical laboratories. Table VI sets forth an analysis of several such kits, and a comparison of the types of protein found in those kits with hog IF, hog R and hog IFC. By reference to Table VI,
35 it is seen that the commercial kits appear to have only about 13% to about 35% intrinsic factor and from about 60% to about 85% R.protein. It is therefore suspected, that the use of these kits will give substantially erroneous assays of the amount of vitamin B₁₂ present in a sample



Table VI
Analysis of vitamin B₁₂ binders and assay pH used in commercial kits sold for the assay of vitamin B₁₂ in clinical laboratories

Source of Binder	Vitamin B ₁₂ Binding Protein		Assay pH of Kit(d)
	intrinsic factor(a) (%)	R protein(b) (%)	
Hog IF	97	0	-
Hog R	0	98	-
Hog IFC	25	75	-
Diagnostic Products Corp. Kit	35	60	1.7
New England Nuclear Kit	20	82	4.1
Bio-Rad Laboratories Kit	30	71	1.9
Medvak Diagnostic Products Kit	13	85	1.8
Schwarz/Mann Kit	34	67	9.1
Pharmacia Diagnostics Kit	1 (33) ^c	67	4.1

- (a) Inhibition of [⁵⁷Co] B₁₂ binding observed with anti-IF antibodies at pH 7.5.
- (b) Inhibition of [⁵⁷Co] B₁₂ binding observed with a 100 fold molar excess of cobinamide at pH 7.5.
- (c) In this kit the binder is covalently attached to an insoluble matrix and because of steric factors it may not be accessible to anti-intrinsic factor antibodies. Thus the value for % intrinsic factor may be as high as 33%.
- (d) Refers to the pH measured after the addition of all of the components of each individual kit. R protein retains its full B₁₂ binding ability over the range of pH 1.7-9.1 but intrinsic factor loses 10% and 99% of its B₁₂ binding ability at pH 4.1 and 1.9, respectively. Thus kits that use pH 4.1 have slightly less % intrinsic factor than is indicated in the table while those that use pH 1.7-1.9 are employing essentially no intrinsic factor.



when the sample also includes vitamin B₁₂ analogues, such as those newly discovered to exist in mammalian blood and tissues. It has also been determined that the effectiveness of intrinsic factor to bind vitamin B₁₂ is somewhat
5 pH dependent, with intrinsic factor losing about 10% of its binding ability at a pH of about 4.1 and losing about 99% of its binding ability at a pH of about 1.9. Thus, to the extent that the commercial kits use a pH of about 4.1 during binding they would have about 10% less intrinsic
10 factor than shown in Table VI. Those kits having a pH during binding of about 1.7 to about 1.9 would obtain substantially no binding from intrinsic factor.

While the invention has been particularly shown and described with reference to preferred embodiments thereof,
15 it will be understood by those skilled in the art that the foregoing and other modifications or changes in form and details may be made therein without departing from the spirit and scope of the invention.

What is claimed is:



CLAIMS

32

1. A composition for use in the assay of vitamin B₁₂ (cobalamin) wherein the active component is substantially specific to vitamin B₁₂ and substantially free of substances which react with vitamin B₁₂ analogues.
2. The composition of claim 1 wherein the active component is a binding protein.
3. The composition of claim 2 wherein the binding protein is intrinsic factor.
4. The composition of claim 3 wherein the intrinsic factor is selected from the group consisting of human IF, hog IF and rabbit IF.
5. The composition of claim 4 wherein the intrinsic factor includes human IF.
6. The composition of claim 4 wherein the intrinsic factor includes hog IF.
7. The composition of claim 1 wherein one or more protein binder which is non-specific to vitamin B₁₂ is present and has been rendered substantially inactive or inoperative as to its ability to bind with vitamin B₁₂ by treatment with a vitamin B₁₂ analogue, said analogue being substantially reactive with said non-specific protein and substantially non-reactive with the material which is a specific binder for B₁₂, said analogue being present in an amount sufficient to render substantially all of the non-specific binder inactive or inoperative in its ability to bind with vitamin B₁₂.
8. The composition of claim 7 wherein the specific active binder includes intrinsic factor protein and the non-specific protein binder includes an R protein.
9. The composition of claim 8 wherein a portion of the vitamin B₁₂ analogue is selected from the group consisting of cobinamide, CN-Cbl(bde-OH) and (3,5,6-Me₃BZA) (CN,OH) Cba.

BEST AVAILABLE COPY



10. The composition of claim 7 wherein the vitamin B₁₂ analogue is present in the amount of about a 10-fold to about a 100,000-fold excess, based on the molar amount of the non-specific protein.
11. The composition of claim 9 wherein the vitamin B₁₂ analogue is present in the amount of about a 1-fold to about a 10,000,000-fold excess based on the molar amount of the non-specific protein.
12. The method of preparing the binder of claim 2 wherein a composition including binding material substantially specific to vitamin B₁₂ and binding materials substantially non-specific to vitamin B₁₂ and including a protein is treated with vitamin B₁₂ analogues to remove substantially all of the non-specific vitamin B₁₂ binding material, said vitamin B₁₂ analogue being substantially reactive with said non-specific binder and substantially non-reactive with the material which is a specific binder for B₁₂, said analogue being added in an amount sufficient to render substantially all of the non-specific binder inactive or inoperative in its ability to bind with vitamin B₁₂.
13. The method of preparing the binder of claim 7 wherein a composition including protein binding material substantially specific to vitamin B₁₂ and binding material substantially non-specific to vitamin B₁₂ is treated with vitamin B₁₂ analogue to render the binder non-specific to vitamin B₁₂ substantially inactive or inoperative as to its ability to bind with vitamin B₁₂, said vitamin B₁₂ analogue being substantially reactive with said non-specific binder and substantially non-reactive with the material which is a specific binder for B₁₂, said analogue being added in an amount sufficient to render substantially all of the non-specific binder inactive or inoperative in its ability to bind with vitamin B₁₂.

14. The method of claim 13 wherein the specific active binder includes intrinsic factor protein and the non-specific protein binder includes an R protein.

15. A radioisotope dilution assay for measuring the vitamin B₁₂ level in a sample comprising:
contacting said sample with a known amount of a radioisotope of vitamin B₁₂ and a composition containing a binding protein substantially specific to vitamin B₁₂, said composition being substantially free of substances which bind vitamin B₁₂ analogues.

16. The method of claim 15 wherein the binding protein is intrinsic factor.

17. The method of claim 15 wherein the binding protein consists essentially of protein selected from the group consisting of human IF, hog IF and rabbit IF.

18. The method of claim 15 wherein the source of vitamin B₁₂ is selected from the group consisting of mammalian tissue and mammalian blood.

19. The method of claim 15 wherein the source of vitamin B₁₂ is human blood.

20. The method of measuring vitamin B₁₂ analogues present in a sample which also includes vitamin B₁₂, including the steps of:
assaying a portion of the sample with a binder which is a binder for both vitamin B₁₂ and vitamin B₁₂ analogues;

assaying a portion of the sample with a binder which is substantially specific to vitamin B₁₂ and substantially free of substances which bind vitamin B₁₂ analogues; and

then determining the difference between the two assays as being substantially indicative of the amount of vitamin B₁₂ analogue in the sample.

BAD ORIGINAL

BEST AVAILABLE COPY



INTERNATIONAL SEARCH REPORT

0880047 QW

International Application No PCT/US79/00210

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC INT. CL. GOIN 33/16; U.S. CL. 23/230B, 230.3, 230.6; 252/408; 424/1.1, 5, 96		
II. FIELDS SEARCHED		
Minimum Documentation Searched *		
Classification System: US Classification Symbols: 23/230B, 230.3, 230.6, 195/2, 252/408, 424/1.1, 5, 96		
US	23/230B, 230.3, 230.6, 195/2, 252/408, 424/1.1, 5, 96	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched *		
III. DOCUMENTS CONSIDERED TO BE RELEVANT *		
Category *	Citation of Document, ¹⁶ with Indication, where appropriate, of the relevant passages ¹⁷	Relevant to Claim No. ¹⁸
X	US, A, 3,434,927 Published 25 March 1969, Highley et al.	1-11
X	US, A, 3,442,819 Published 06 May 1969, See Examples I-III, Herbert.	15-20
X	US, A, 3,937,799 Published 10 Feb. 1976, Lewis et al.	15-20
X	US, A, 3,591,678 Published 06 July 1971, Ellenbogen et al.	1-11
A	US, A, 2,892,754 Published 30 June 1959, Nimeguen.	1-11
A	US, A, 2,912,360 Published 10 Nov. 1959, Baum et al.	1-11
A	US, A, 3,980,764 Published 14 Sept. 1976, Adams.	1-11
A, P	US, A, 4,133,951 Published 09 Jan. 1979, Charlton et al.	1-11
A, P	US, A, 4,136,159 Published 23 Jan. 1979, Stone.	1-11
A, P	US, A, 4,146,602 Published 27 March 1979, Gutcho et al.	1-11
A	GB, A, 771,068 Published 27 March 1957.	1-11
A	GB, A, 791,186 Published 26 Feb. 1958.	1-11
A	GB, A, 900,459 Published 04 July 1962.	1-11
<p>* Special categories of cited documents: ¹⁵</p> <p>"A" document defining the general state of the art</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document cited for special reason other than those referred to in the other categories</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but on or after the priority date claimed</p> <p>"T" later document published on or after the international filing date or priority date and not in conflict with the application, but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search *		Date of Mailing of this International Search Report *
10 July 1979		17 JUL 1979
International Searching Authority *		Signature of Authorized Officer ²⁰
ISA/US		Sidney Marantz SIDNEY MARANTZ

Form PCT/ISA/210 (second sheet) (October 1977)

BAD ORIGINAL

BEST AVAILABLE COPY